



Continental-Scale Governance and the Hastening of Loss of Australia's Biodiversity

Against a global backdrop of rapid environmental change, conserving biodiversity poses one of the biggest and most important challenges to society. For this reason, systems of nature reserves have never been more important. Protected areas are under threat in many parts of the world (Mascia & Pailler 2011), but the weakening of protected areas in a rich, developed country with a global reputation for conservation leadership (Harrison 2006) is particularly alarming (Ritchie 2013). Consequently, we are concerned about the recent spate of substantial policy, legislative, and management changes being made by three of six Australian state governments for exploitative uses of national parks—actions that could affect much of Australia and have negative effects on biodiversity.

In recent decades, the Australian state and federal governments have collectively built a system of terrestrial and marine conservation reserves that forms the cornerstone of biodiversity conservation on the continent and aspires to be comprehensive and adequate. The resulting national reserve system is imperfect, but it goes some way toward protecting Australia's unique species and ecosystems (Taylor et al. 2011). That system is now being systematically undermined, even though continental-scale biodiversity losses are underway.

Australia's highly diverse and predominantly endemic biodiversity is seriously imperiled. In the past 200 years, at least 27 mammals, 23 birds (including island species and subspecies), 4 frogs, and over 60 plant species have become extinct (Department of Sustainability 2009). In addition, over 1500 mammals, birds, reptiles, amphibians, and plants and over 3000 ecosystem types are currently threatened with extinction (Keith et al. 2013). In Victoria, for instance, only approximately 30% of the original native vegetation remains, and some vegetation types, such as grasslands and open woodlands, have been reduced by more than 99% since European settlement (Bradshaw 2012). The situation for marine systems is far more uncertain owing to data limitations even for economically important species (Beeton et al. 2012; FRDC 2012). In addition, Australia has the world's most recent mammal extinction, the Christmas Island pipistrelle bat (*Pipistrellus murrayi*) in 2009 (Martin et al. 2012). If current trends continue, many other species such as the Leadbeater's possum (*Gymnobelideus leadbeateri*)

will have the same fate. Lindenmayer and Possingham (2013) suggest the Victorian government is knowingly condoning activities that will reduce the viability of this species, which is listed by the International Union for Conservation of Nature as endangered.

Given these realities, it is not surprising that proposals to weaken the country's nature reserves are raising alarm bells among conservation biologists and concerned members of the public. Recently proposed or enacted laws will allow an increase in exploitative uses of reserves—including industrial logging, grazing by domestic livestock, mining, commercial development, and recreational hunting and fishing—all of which are detrimental to nature conservation. The overall conservation effect and prognosis worsens because these same Australian state governments are reversing safeguards that curb the clearing of native vegetation outside protected areas. Remnants of many Australian ecosystems persist mostly on private and leasehold land (Benson 2008) or in unreserved marine areas, and these provide a necessary complement to the biodiversity protection offered in reserves. In Queensland and Victoria, hard-won laws constraining vegetation clearing on private land are now being relaxed, and this will certainly accelerate the loss of regional biodiversity. In Queensland these regressive changes add to the already tenuous status of any conservation covenants with potential for mining exploration and development (Adams & Moon 2013). In Western Australia there have been large excisions of existing conservation land for mining. The government of New South Wales is considering relaxing anticlearing laws, even though Pressey et al. (2000) demonstrated that 85% of the state's native vegetation with high conservation priority was on private land. Just as troubling, legal-aid funding in New South Wales is being wound back for public-interest environmental cases, making it even more difficult to bring the state government to environmental account (Smith 2013).

Collectively, these new proposals represent a serious about-face for government policy on nature conservation in Australia. They will increase the dependence of Australian biodiversity on protected areas and reduce reserve viability by weakening biodiversity protections inside reserves and reducing ecological connectivity and

accelerating habitat loss outside reserves. Laurance et al. (2012) found that pressures outside reserves negatively affect species residing within them. Species are likely to be further affected by a disrupted climate and by increasing pressures from invasive pests, fire, disease, and drought (Brook et al. 2008). Even before these changes, Australia's reserve network was showing signs of inadequacy, and there were documented collapses in regional faunal communities within national parks (Mac Nally et al. 2009; Woinarski et al. 2011).

Economic rationales are being used to justify the dismantling of park protections. These arguments include providing opportunities to feed cattle in national parks during drought and assisting ailing forestry industries by opening up new areas for logging. However, the use of reserves for such activities has substantial long-term costs. Additional environmental impacts include a likely increase in weed establishment, higher carbon emissions, lower carbon sequestration capacity, increased soil erosion, and damage to sensitive riparian zones and waterways. Indeed, the effect of cattle grazing on ecosystems is likely to be most severe during drought because the effects of drought and grazing can interact (Loeser et al. 2007). The repair bill for these effects will dwarf any short-term economic benefits to extractive industries, and some changes might be irreversible (Cardinale et al. 2012). The fact that state governments are retreating from the previously accepted principal purpose of reserves—to conserve biodiversity—suggests a shortsighted decline in political and societal concern for nature conservation (McCallum & Bury 2013).

Hunting and fishing lobbies are also arguing for increased access to nature reserves. One argument suggests these activities could help control problem species. Hunting to manage pest species can be justified occasionally in national parks, but it must be carefully based on scientific evidence. One serious possible consequence of nonstrategic hunting is that removing one nonnative species without considering its interactions with and effect on other species could usher in unexpected and disadvantageous ecological consequences (e.g., increase abundances of some pest species following release from another competitor or predator and subsequent higher rates of predation on their native prey) (Ritchie & Johnson 2009; Ruscoe et al. 2011). Of course, hunting might have no effect on the target species. For most species, effective population reduction through hunting seems unlikely due to factors such as the cryptic behavior and patchy distribution, often intrinsically high rates of increase, and compensatory breeding or survival of invasive nonnative animals and the vast, remote, and often rugged nature of many of Australia's national parks, which makes widespread access for hunting difficult (Booth 2010). Other forms of pest control that involve encouraging or reintroducing apex predators (e.g., dingoes

[*Canis lupus dingo*]) could be far more effective ecologically and economically in the long term (Ritchie et al. 2012).

Opening up Australia's conservation reserves to extractive uses, whether to support primary industries or to allow hunting and fishing, is all the less defensible because conservation reserves in Australia already occupy the margins of productivity for agriculture, grazing, and logging (Pressey et al. 2000). Although Australia's national reserve system covers 13.4% of the country's landmass (over 500 national parks cover 3.6%), they protect a disproportionately small percentage of productive landscapes. Similarly, marine sanctuary zones represent about 5% of nearshore and 15% of Commonwealth waters (>3 nautical miles offshore) but are strongly biased toward areas with least value to commercial fishing and without value for fossil-fuel development (Barr & Possingham 2013). In other words, protected areas present almost no barrier to economic development in Australia. The biased character of Australia's reserves underlines the importance of stronger, not weaker, protection of biodiversity outside reserves.

The recent legislative threats to Australia's parks have come about not because of a lack of data on their likely consequences (Lindenmayer & Possingham 2013), but apparently to cater to particular political interests. Given that continued weakness in the global economy will increase incentives for governments to allow further exploitation of natural resources, we must remain vigilant to the potentially serious consequences of such actions. There are lessons here for conservation globally: laws for nature conservation can be undone because governments change; even wealthy countries are willing to sacrifice long-term conservation outcomes for the possibility of short-term economic gains (Bradshaw et al. 2010); established conservation reserves might need multiple layers of protection from the vagaries of policy and legislation; and as conservation biologists, we need to work harder to build public constituencies that support the protection of reserves and fight against watering down important environmental legislation.

Poorly framed and opportunistic legislation that will erode the ecological integrity and conservation value of protected areas and off-reserve management needs to be rescinded. The scientific evidence to support the importance to biodiversity of maintaining a well-managed system of protected areas in terrestrial and marine landscapes is overwhelming. Ultimately, the commitment of any government to nature conservation will be measured not by hectares under nominal protection, but by what development potential it is prepared to forgo to avoid the loss of biodiversity. Although Australia's nominally protected areas increase in area, the trajectory of real commitment to conservation is in decline along with Australia's biodiversity.

Euan G. Ritchie,* Corey J. A. Bradshaw,†
Chris R. Dickman,‡ Richard Hobbs,§
Christopher N. Johnson, Emma L. Johnston,††**
William F. Laurance,‡‡ David Lindenmayer,§§
Michael A. McCarthy, Dale G. Nimmo,***
Hugh H. Possingham,††† Robert L. Pressey,‡‡‡
David M. Watson,§§§ and John Woinarski****

*Centre for Integrative Ecology and School of Life and Environmental Sciences, Deakin University, Burwood, Victoria 3125, Australia email e.ritchie@deakin.edu.au

†The Environment Institute and School of Earth and Environmental Sciences, The University of Adelaide, Adelaide, South Australia 5005, Australia

‡School of Biological Sciences, University of Sydney, New South Wales 2006, Australia

§School of Plant Biology, University of Western Australia, Western Australia 6009, Australia

**School of Zoology, University of Tasmania, Hobart, Tasmania 7001, Australia

††Evolution and Ecology Research Centre, University of New South Wales, Sydney, New South Wales 2052, Australia

‡‡Centre for Tropical Environmental and Sustainability Science and School of Marine and Tropical Biology, James Cook University, Cairns, Queensland 4870, Australia

§§Australian Research Council Centre of Excellence for Environmental Decisions Fenner School of Environment and Society, Australian National University, Canberra, Australian Capital Territory 0200, Australia

***School of Botany, The University of Melbourne, Parkville, Victoria 3010, Australia

†††Australian Research Council Centre of Excellence for Environmental Decisions, The University of Queensland, Brisbane, Queensland 4072, Australia

‡‡‡Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland 4811, Australia

§§§Institute for Land, Water & Society, Charles Sturt University, Albury, New South Wales 2640, Australia

****Research Institute for the Environment and Livelihoods, Charles Darwin University, Casuarina, Northern Territory 0909, Australia

Literature Cited

- Adams, V. M., and K. Moon. 2013. Security and equity of conservation covenants: contradictions of private protected area policies in Australia. *Land Use Policy* **30**:114–119.
- Barr, L. M., and H. P. Possingham. 2013. Are outcomes matching policy commitments in Australian marine conservation planning? *Marine Policy* **42**:39–48.
- Beeton, R. J. S., C. D. Buxton, G. C. Cutbush, P. G. Fairweather, E. L. Johnston, and R. Ryan. 2012. Report of the Independent Scientific Audit of Marine Parks in New South Wales. NSW Department of Primary Industries and Office of Environment and Heritage, NSW (p. 1–124).
- Benson, J. S. 2008. NSW Vegetation classification and Assessment: part 2 plant communities of the NSW South-western Slopes Bioregion and update of NSW Western Plains plant communities. Version 2 of the NSWVCA database. *Cunninghamia* **10**:599–673.
- Booth, C. 2010. Hunting and feral animal control: Conservation or con? Pages 25–31 in M. Tensen and B. Jones, editors. *Convergence or conflict: animal welfare in wildlife management and conservation*. Royal Society for the Prevention of Cruelty to Animals, Canberra.
- Bradshaw, C. J. A. 2012. Little left to lose: deforestation and forest degradation in Australia since European colonization. *Journal of Plant Ecology* **5**:109–120.
- Bradshaw, C. J. A., X. Giam, and N. S. Sodhi. 2010. Evaluating the relative environmental impact of countries. *PLoS ONE* **5**: DOI: 10.1371/journal.pone.0010440
- Brook, B. W., N. S. Sodhi, and C. J. A. Bradshaw. 2008. Synergies among extinction drivers under global change. *Trends in Ecology & Evolution* **23**:453–460.
- Cardinale, B. J., et al. 2012. Biodiversity loss and its impact on humanity. *Nature* **486**:59–67.
- Department of Sustainability, Environment, Water, Population and Communities. 2009. EPBC Act list of threatened fauna. Australian Government, Canberra. Available from <http://www.environment.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl?wanted=faua> (accessed August 2013).
- Fisheries Research & Development Corporation. 2012. Status of key Australian fish stocks. FRDC, Canberra.
- Harrison, A. L. 2006. Who's who in *Conservation Biology*—an authorship analysis. *Conservation Biology* **20**:652–657.
- Keith, D. A., et al. 2013. Scientific foundations for an IUCN Red List of ecosystems. *PLoS ONE* **8** DOI: 10.1371/journal.pone.0062111
- Laurance, W. F., et al. 2012. Averting biodiversity collapse in tropical forest protected areas. *Nature* **489**:290–294.
- Lindenmayer, D. B., and H. P. Possingham. 2013. No excuse for habitat destruction. *Science* **340**:680.
- Loeser, M. R. R., T. D. Sisk, and T. E. Crews. 2007. Impact of grazing intensity during drought in an Arizona grassland: Contributed papers. *Conservation Biology* **21**:87–97.
- Mac Nally, R., A. F. Bennett, J. R. Thomson, J. Q. Radford, G. Unmack, G. Horrocks, and P. A. Vesk. 2009. Collapse of an avifauna: climate change appears to exacerbate habitat loss and degradation. *Diversity and Distributions* **15**:720–730.
- Martin, T. G., S. Nally, A. A. Burbidge, S. Arnall, S. T. Garnett, M. W. Hayward, L. F. Lumsden, P. Menkhorst, E. McDonald-Madden, and H. P. Possingham. 2012. Acting fast helps avoid extinction. *Conservation Letters* **5**:274–280.
- Mascia, M. B., and S. Pailler. 2011. Protected area downgrading, downsizing, and degazettement (PADDD) and its conservation implications. *Conservation Letters* **4**:9–20.
- McCallum, M., and G. Bury. 2013. Google search patterns suggest declining interest in the environment. *Biodiversity and Conservation* **22**:1355–1367.
- Pressey, R. L., T. C. Hager, K. M. Ryan, J. Schwarz, S. Wall, S. Ferrier, and P. M. Creaser. 2000. Using abiotic data for conservation assessments over extensive regions: quantitative methods applied across New South Wales, Australia. *Biological Conservation* **96**:55–82.
- Ritchie, E. G. 2013. Conservation: relaxed laws imperil Australian wildlife. *Nature* **498**:434.
- Ritchie, E. G., B. Elmhagen, A. S. Glen, M. Letnic, G. Ludwig, and R. A. McDonald. 2012. Ecosystem restoration with teeth: What role for predators? *Trends in Ecology & Evolution* **27**:265–271.
- Ritchie, E. G., and C. N. Johnson. 2009. Predator interactions, meso-predator release and biodiversity conservation. *Ecology Letters* **12**:982–998.
- Ruscoe, W. A., et al. 2011. Unexpected consequences of control: competitive vs. predator release in a four-species assemblage of invasive mammals. *Ecology Letters* **14**:1035–1042.
- Smith, J. 2013. Legal aid cuts threaten environmental justice. *EDO NSW Weekly Bulletin* **814**:1.
- Taylor, M. F. J., P. S. Sattler, M. Evans, R. A. Fuller, J. E. M. Watson, and H. P. Possingham. 2011. What works for threatened species recovery? An empirical evaluation for Australia. *Biodiversity and Conservation* **20**:767–777.
- Woinarski, J. C. Z., et al. 2011. The disappearing mammal fauna of northern Australia: context, cause, and response. *Conservation Letters* **4**:192–201.